

Laplacian eigenmodes for spherical spaces

M. Lachièze-Rey ^{1,2} and S. Caillerie ¹

1. Service d'Astrophysique, C. E. Saclay

91191 Gif sur Yvette cedex, France

2. CNRS : UMR 7164

February 2, 2008

Abstract

The possibility that our space is multi - rather than singly - connected has gained a renewed interest after the discovery of the low power for the first multipoles of the CMB by WMAP. To test the possibility that our space is a multi-connected spherical space, it is necessary to know the eigenmodes of such spaces. Excepted for lens and prism space, and in some extent for dodecahedral space, this remains an open problem. Here we derive the eigenmodes of all spherical spaces. For dodecahedral space, the demonstration is much shorter, and the calculation method much simpler than before. We also apply to tetrahedric, octahedric and icosahedric spaces. This completes the knowledge of eigenmodes for spherical spaces, and opens the door to new observational tests of cosmic topology.

The vector space $\mathcal{V}^k$ of the eigenfunctions of the Laplacian on the three-sphere $\mathcal{S}^3$, corresponding to the same eigenvalue $\lambda_k = -k(k+2)$, has dimension $(k+1)^2$. We show that the Wigner functions provide a basis for such space. Using the properties of the latter, we express the behavior of a general function of $\mathcal{V}^k$ under an arbitrary rotation G of $\text{SO}(4)$. This offers the possibility to select those functions of $\mathcal{V}^k$ which remain invariant under G .

Specifying G to be a generator of the holonomy group of a spherical space X , we give the expression of the vector space $\mathcal{V}_X^k$ of the eigenfunctions of X . We provide a method to calculate the eigenmodes up to arbitrary order. As an illustration, we give the first modes for the spherical spaces mentioned.

1 Introduction

1.1 Cosmic Topology

The possibility that spatial sections of our Universe have a multi-connected topology offers interesting alternatives to the standard big bang models [10]. The recent suggestion [13] that a peculiar variant, namely the Poincaré dodecahedral space, may explain the deficiency in the first modes of the angular power spectrum of the CMB, as observed by WMAP has renewed the interest to this question. It is presently an important task of cosmology to check this possibility.

The best hope to test the possible multi-connectedness of space comes from the CMB fluctuations. For this task, the most important characteristic of a multi-connected space X is the set of its eigenmodes, a subspace of the modes of its universal covering [15]. The main consequences on the CMB are (i) a suppression of power at the scales comparable to, or larger than, the circumference of X , and (ii) a violation of isotropy at these scales. Thus, observational signatures may occur as a reduction of power in the angular CMB spectrum (as reported by [13]), and as the imprint of correlations between different modes. The predictions of these observable effects require the knowledge of the eigenmodes [of the Laplacian] of X . For instance, [13] use a limited number of eigenmodes of the Poincaré space, calculated numerically. More precise checks require the knowledge of a larger number of modes of this space. The examination of other models require the knowledge of the eigenmodes of other spherical spaces.

This paper is devoted to the calculation of the eigenmodes of all variants of the multi-connected spherical spaces. They are defined as quotients $X \equiv \mathcal{S}^3/\Gamma$, where the three-sphere $\mathcal{S}^3$ is the universal covering space of X , and $\Gamma \subset \text{SO}(4)$ is the holonomy group of X . The transformations of Γ leave X invariant. The compactness of $\mathcal{S}^3$ implies the finiteness of Γ . The latter is finitely generated, *i.e.* by a finite number of transformations.

1.2 Spherical spaces

In the family of spherical spaces, the eigenmodes of lens and prism spaces have been calculated by [11], [12] (see also [8]) in analytic form. For the dodecahedral space, only the very first modes had been calculated numerically, by [13]. Later, [9] reduced the calculation to an eigenvalue problem of reduced dimensionality, allowing their estimations up to arbitrary order. For the other spherical spaces, the knowledge of the eigenmodes remains an open problem. Here, we present a general method to calculate the eigenmodes of any of the spherical spaces $X = \mathcal{S}^3/\Gamma$.

For a given eigenvalue λ_k , the eigenmodes of X describe an eigenspace $\mathcal{V}_X^k$. The dimension of this vector space has been calculated, very generally, by [6]. We use his results as a cross-check of the calculations below.

Our calculation is based on the fact that each Γ has two generators $G_\pm$, which are single left action rotations of $\text{SO}(4)$. We obtain the straightforward decomposition

$$\mathcal{V}_X^k = \mathcal{V}_{G_+}^k \cap \mathcal{V}_{G_-}^k,$$

and we give explicit analytic expressions of the $\mathcal{V}_{G_\pm}^k$, which are the vector spaces of the eigenfunctions of $\mathcal{S}^3$ invariant under $G_\pm$.

Although the expressions of the two $\mathcal{V}_{G_\pm}^k$'s are very simple, there is no general analytic expression of their intersection. Thus, to obtain an explicit expression of the modes, we propose a general method, which applies to any spherical space. It is based on the fact that a basis of $\mathcal{V}^k$ is provided by the Wigner functions. From the rotation properties of the latter, we derive a simple invariance condition of the modes of $\mathcal{S}^3$ under $G_\pm$, and thus under Γ . This provides the eigenmodes of X , which are the eigenmodes of $\mathcal{S}^3$ which remain invariant under Γ .

In (2), we analyze the eigenfunctions of $\mathcal{S}^3$. We prove that the Wigner functions provide a convenient basis ($\mathcal{B}_2$), different from the widely used basis ($\mathcal{B}_1$) provided by the usual hyperspherical harmonics. In (3), we derive the rotation properties of this basis under $\text{SO}(4)$, from which we

deduce those of all eigenfunctions of $\mathcal{S}^3$. This allows to give a criterion for the invariance of such a function under an arbitrary rotation of $\text{SO}(4)$. In (4), we give an explicit analytic expression for the invariant subspaces $\mathcal{V}_{G_{\pm}}^k$, and we provide a general procedure to calculate the eigenmodes of any spherical space. The first modes are explicitly given in Appendix.

2 Eigenmodes of $\mathcal{S}^3$

The eigenmodes of $\mathcal{S}^3$ are the eigenfunctions of the Laplacian $\Delta_{\mathcal{S}^3}$ of $\mathcal{S}^3$. The eigenvalues are known to be of the form $\lambda_k = -k(k+2)$, where $k \in \mathbb{N}$. The eigenmodes corresponding to a given value of k span the eigenspace $\mathcal{V}^k$, a vector space of dimension $(k+1)^2$. They are the solutions of the Helmholtz equation

$$\Delta_{\mathcal{S}^3} f = \lambda_k f = -k(k+2) f, \quad k \in \mathbb{N}. \quad (1)$$

Each eigenspace $\mathcal{V}^k$ realizes the $(k+1)^2$ dimensional irreducible representation of $\text{SO}(4)$, the isometry group of $\mathcal{S}^3$.

The corresponding eigenspace $\mathcal{V}_X^k$, of $X = \mathcal{S}^3/\Gamma$, the goal of our search, is the vector space of functions of $\mathcal{V}^k$ which are Γ -invariant. Each $\mathcal{V}_X^k$ is a subvector space of $\mathcal{V}^k$, whose dimension (possibly zero) is the degeneracy of λ_k on X .

2.1 Basis $(\mathcal{B}_1)$: hyperspherical harmonics

Let us call $(\mathcal{B}_1)$ the most widely used (orthonormal) basis for $\mathcal{V}^k$ provided by the hyperspherical harmonics

$$(\mathcal{B}_1) \equiv \{\mathcal{Y}_{k\ell m}, \quad 0 \leq \ell \leq k, -\ell \leq m \leq \ell\} \quad (2)$$

It generalizes the usual spherical harmonics $Y_{\ell m}$ on the sphere (note that $\mathcal{Y}_{k\ell m} \propto Y_{\ell m}$). In fact, it can be shown ([1], [3] p.240,[4]) that a basis of this type exists on any sphere S^n . Moreover, [3] [4] show that the $(\mathcal{B}_1)$ basis for S^n is “naturally generated” by the $(\mathcal{B}_1)$ basis for S^{n-1} . In this sense, the $(\mathcal{B}_1)$ basis for $\mathcal{S}^3$ is generated by the usual spherical harmonics $Y_{\ell m}$ on the 2-sphere S^2 .

The generation process involves harmonic polynomials constructed from null complex vectors. The basis $(\mathcal{B}_1)$ is in fact based on the reduction of the representation of $\text{SO}(4)$ to representations of $\text{SO}(3)$: each $\mathcal{Y}_{k\ell m}$ is an eigenfunction of an $\text{SO}(3)$ subgroup of $\text{SO}(4)$ which leaves a selected point of $\mathcal{S}^3$ invariant. This makes these functions useful when one considers the action of that particular $\text{SO}(3)$ subgroup. But they show no simple behaviour under a general rotation.

This basis is adapted to the usual polar spherical coordinates (ψ, θ, ϕ) .

2.2 The parabolic basis $(\mathcal{B}_2)$

The second (orthonormal) basis of $\mathcal{V}^k$ is specific to $\mathcal{S}^3$:

$$(\mathcal{B}_2) \equiv \{T_{k;m_1, m_2}, \quad -k/2 \leq m_1, m_2 \leq k/2\} \quad (3)$$

where m_1 and m_2 vary independently by integers (and, thus, take integral or semi-integral values according to the parity of k). It is for instance

introduced in [1] by group theoretical arguments, it appears naturally adapted to the systems of toroidal coordinates to describe $\mathcal{S}^3$:

$$\begin{cases} x^0 &= r \cos \chi \cos \theta \\ x^1 &= r \sin \chi \cos \phi \\ x^2 &= r \sin \chi \sin \phi \\ x^3 &= r \cos \chi \sin \theta \end{cases}$$

spanning the range $0 \leq \chi \leq \pi/2$, $0 \leq \theta \leq 2\pi$ and $0 \leq \phi \leq 2\pi$. Here, the (x^μ) represent the cartesian coordinates of the embedding space $\mathbb{R}^4$ (see [8], [11]), and as we consider only spherical spaces, r is thus the radius of $\mathcal{S}^3$, which is a constant assumed hereafter to be 1.

The explicit expression of the $T_{k;m_1,m_2}$ can be found in ([1], [11], see [8]). It is proportional to a Jacobi polynomial:

$$T_{k;m_1,m_2}(X) = C_{k;m_1,m_2} (\cos \chi e^{i\theta})^\ell (\sin \chi e^{i\phi})^m P_d^{(m,\ell)}(\cos(2\chi)), \quad (4)$$

where we have written

$$\ell = m_1 + m_2, \quad m = m_2 - m_1, \quad d = k/2 - m_2.$$

The factor $C_{k;m_1,m_2}$ can be computed from normalization requirements (there is a factor $\sqrt{2}$ lacking in [8] formula) as

$$C_{k;m_1,m_2} \equiv \sqrt{\frac{k+1}{2\pi^2}} \sqrt{\frac{(k/2+m_2)!(k/2-m_2)!}{(k/2+m_1)!(k/2-m_1)!}}. \quad (5)$$

We also note these useful proportionality relations:

$$\begin{aligned} \cos^\ell \chi \sin^m \chi P_{\frac{k-\ell-m}{2}}^{(m,\ell)}(\cos 2\chi) &\propto \cos^\ell \chi \sin^{-m} \chi P_{\frac{k-\ell+m}{2}}^{(-m,\ell)}(\cos 2\chi) \\ &\propto \cos^{-\ell} \chi \sin^m \chi P_{\frac{k+\ell-m}{2}}^{(m,-\ell)}(\cos 2\chi). \end{aligned} \quad (6)$$

Since $(\mathcal{B}_2)$ is a basis, we have $\mathcal{V}^k = \text{span}(T_{k;m_1,m_2})_{-k/2 \leq m_1, m_2 \leq k/2}$. It appears convenient to define the subvector spaces

$$\mathcal{V}^{k,m_1} \equiv \text{span}(T_{k;m_1,m_2})_{-k/2 \leq m_2 \leq k/2}. \quad (7)$$

We have

$$\mathcal{V}^k = \bigoplus_{m_1=-k/2}^{k/2} \mathcal{V}^{k,m_1}.$$

We show now that the $T_{k;m_1,m_2}$ identify, up to a constant, to the Wigner D-functions.

2.3 The Wigner D-functions

The Wigner D-functions are defined as functions on the Lie group $\text{SO}(3)$:

$$\begin{aligned} \text{SO}(3) &\longrightarrow \mathbb{C} \\ g &\longmapsto \mathcal{D}_{m'm}^j(g) \equiv \langle jm' | R_g | jm \rangle. \end{aligned} \quad (8)$$

Here, g is rotation of $\text{SO}(3)$, and R_g expresses its natural left action $f \mapsto R_g f$ on functions: $R_g f(x) \equiv f(g^{-1}x)$. The vectors $|jm\rangle$, $m \in \llbracket -j, j \rrbracket$ form an orthonormal basis for $\mathcal{H}^j$, the $2j+1$ dimensional irreducible representation of $\text{SU}(2)$. When j is integer, this is an IUR of $\text{SO}(3)$, and

the $|jm\rangle$ can be taken as the usual spherical harmonics $Y_{\ell m}$. This implies

$$R_g Y_{jm} = \sum_{m'} \mathcal{D}_{m'm}^j(g^{-1}) Y_{jm'} \quad (9)$$

(Care must be taken that one finds in the literature other definitions with indices exchanged; we adopt here the notations of [2]).

As manifolds, the three-sphere $\mathcal{S}^3$ and $SU(2)$ are identical. This allows to identify any point of $\mathcal{S}^3$ to an element of $SU(2)$ by the following relation:

$$\begin{aligned} \mathcal{S}^3 &\longrightarrow SU(2) \\ x \equiv (\chi, \theta, \phi) &\longmapsto u_x \equiv \begin{pmatrix} \cos \chi & e^{i\theta} & i \sin \chi & e^{i\phi} \\ i \sin \chi & e^{-i\phi} & \cos \chi & e^{-i\theta} \end{pmatrix}. \end{aligned} \quad (10)$$

(Note that one finds other phase choices for this identification in the literature).

On the other hand, there is a group isomorphism between $SU(2)$ and $SO(3)/\mathbb{Z}_2$, where $\mathbb{Z}_2$ refers to the multiplicative group $\{-1, 1\}$. Thus, each element u of $SU(2)$ defines an $SO(3)$ rotation g_u . In practice, a rotation of $SO(3)$ is parametrized by its Euler angles α, β, γ . Taking into account the identification above, the correspondence takes the form:

$$SU(2) \mapsto SO(3) \quad (11)$$

$$u = (\chi, \theta, \phi) \mapsto g_u = (\alpha, \beta, \gamma), \quad (12)$$

where

$$\chi = \frac{\beta}{2} \quad \theta = \frac{\alpha + \gamma}{2} \quad \phi = \frac{\alpha - \gamma}{2}. \quad (13)$$

This allows to consider the Wigner D-functions as functions on $SU(2)$ and, thus, on $\mathcal{S}^3$. Their explicit expression (see for instance [2]) shows their identification to the previous functions $T_{k;m_1,m_2}$:

$$D_{m_2,m_1}^{k/2}(u_x) \equiv \sqrt{\frac{2\pi^2}{k+1}} T_{k;m_1,m_2}(u), \quad (14)$$

with $-k/2 \leq m_1, m_2 \leq k/2$. This will allow to use their very convenient properties with respect to rotations.

2.4 Change of basis

We have two bases for $\mathcal{V}^k$. Calculations may be more convenient in one or the other. Most numerical codes use a development in the basis $(\mathcal{B}_1)$. Here we will calculate the eigenmodes of X in the basis $(\mathcal{B}_2)$. A conversion requires the formulae for the change of basis, which result from the properties of the Wigner functions.

The expansion may be found in [5] (see also [1], which uses however different conventions):

$$\mathcal{Y}_{klm}(u) = i^{k+l-m} \sum_{m_1,m_2} (-1)^{m_2} \left(\frac{k}{2}, -m_1; \frac{k}{2}, m_2 \mid l, m\right) T_{k;m_1,m_2}(u), \quad (15)$$

where the $(j, m_1; j, m_2 \mid l, m)$ are the Clebsch-Gordan coefficients. We can invert this formula by multiplying it by another Clebsch-Gordan coefficient $(j, m_3; j, m_4 \mid l, m)$ and by summing on the 2 indices l and m . The

orthonormality of the Clebsch-Gordan coefficients then gives the relation (after renaming indices):

$$T_{k;m_1,m_2}(u) = (-1)^{m_2} \sum_{l,m} i^{-k-l+m} \left(\frac{k}{2}, -m_1; \frac{k}{2}, m_2 \mid l, m\right) \mathcal{Y}_{klm}(u) \quad (16)$$

The properties of the Clebsch-Gordan coefficients (non zero only when $-m_1 + m_2 = m$ with the previous notation) reduce these formulae, respectively, to

$$\mathcal{Y}_{klm}(u) = \sum_{m_1} i^{k+l+m_1+m_2} \left(\frac{k}{2}, -m_1; \frac{k}{2}, m_1+m \mid l, m\right) T_{k;m_1,m_1+m}(u). \quad (17)$$

and

$$T_{k;m_1,m_2}(u) = \sum_l \left(\frac{k}{2}, -m_1; \frac{k}{2}, m_2 \mid l, m_2 - m_1\right) \frac{\mathcal{Y}_{kl(m_2-m_1)}(u)}{i^{k+l+m_1+m_2}}. \quad (18)$$

3 SO(4) rotations

3.1 General case

The isometries of $\mathcal{S}^3$ form the group $\text{SO}(4)$, the rotation group of the embedding space $\mathbb{R}^4$. We note $G : x \mapsto Gx$ such an isometry, and define its action on a function f of $\mathcal{S}^3$ as

$$R_G : f \mapsto R_G f; \quad R_G f(x) \equiv f(G^{-1}x).$$

Using the previous identification between $\mathcal{S}^3$ and $\text{SU}(2)$, it is very convenient to use the $\text{SU}(2) \times \text{SU}(2)$ representation of $\text{SO}(4)$. Any rotation G of $\text{SO}(4)$ is represented (modulo a sign) as a pair (u_l, u_r) of two $\text{SU}(2)$ matrices, respectively acting on the left and on the right on u_x , an element of $\text{SU}(2)$:

$$\begin{aligned} (u_l, u_r) : \quad \text{SU}(2) &\longrightarrow \text{SU}(2) \\ u_x &\longmapsto u_l \cdot u_x \cdot u_r = u_{Gx}, \end{aligned} \quad (19)$$

where all matrices belong to $\text{SU}(2)$. (The $\text{SO}(4)$ matrices of the form $(u_l, u_r = u_l^{-1})$ form a specific $\text{SO}(3)$ subgroup).

The resolution of identity in $\mathcal{H}^j$ (or equivalently, the composition of two successive $\text{SO}(3)$ rotations, with $R_{g_1 \circ g_2} = R_{g_2} \circ R_{g_1}$), implies the well known addition formula

$$D_{mm'}^j(uv) = \sum_{m''} D_{mm''}^j(u) D_{m''m'}^j(v). \quad (20)$$

Using (14), this implies

$$T_{k;m'm}(x_{uv}) = \sum_{m''} D_{mm''}^{k/2}(u) T_{k;m'm''}(x_v). \quad (21)$$

Here, x_u is the point of $\mathcal{S}^3$ corresponding to $u \in \text{SU}(2)$. Now we will use the same notation for a function on $\mathcal{S}^3$ and on $\text{SU}(2)$: $f(x_u) = f(u)$.

In these formulae, uv refers to the product of two $\text{SU}(2)$ matrices. We are now able to compute the action of a rotation G of $\text{SO}(4)$ on the basis functions $T_{k;m_1,m_2}$ by using the decomposition (19) of G and by using twice the addition formula (21) for $T_{k;m_1,m_2}$.

Thus, we obtain the rotation properties of the basis $(\mathcal{B}_2)$ of eigenfunctions under the rotation $G \approx (u_l, u_r)$:

$$\begin{aligned} R_G T_{k;m_1, m_2}(v) &\equiv T_{k;m_1, m_2}(u_l^{-1} v u_r^{-1}) \\ &= \sum_m \sum_{m'} D_{m_2 m}^{k/2}(u_l^{-1}) D_{m' m_1}^{k/2}(u_r^{-1}) T_{k; m' m}(v). \end{aligned} \quad (22)$$

This completely specifies the rotation properties of the basis $(\mathcal{B}_2)$ under $\text{SO}(4)$.

3.2 Single action rotations

A special case is a single left action rotation (hereafter SLAR), $G \equiv (u_l, u_r \equiv \mathbb{I}_{\text{SU}(2)})$, where $u_l \in \text{SU}(2)$ defines an $\text{SO}(3)$ rotation g . The previous formula reduces to:

$$\begin{aligned} R_G T_{k; M, m_2}(v) &\equiv T_{k; m_1, m_2}(u_l^{-1} v) \\ &= \sum_m D_{m_2 m}^{k/2}(u_l^{-1}) T_{k; M m}(v). \end{aligned} \quad (23)$$

We note that G preserves the first index $M = m_1$. This means that each $\mathcal{V}^{k, M}$ (see 7) is invariant under G . This implies that the research of eigenfunctions invariant under a SLAR G may be performed in each $\mathcal{V}^{k, M}$ separately. Moreover, since the invariance equation does not depend on $M = m_1$, the solution in one of them will give the solution for all of them. From (23), the invariance condition reads

$$T_{k; M, m_2}(v) = \sum_m D_{m_2 m}^{k/2}(u_l^{-1}) T_{k; M m}(v). \quad (24)$$

3.3 Invariant functions

The eigenfunctions of $X \equiv \mathcal{S}^3/\Gamma$ are the eigenfunctions of $\mathcal{S}^3$ which remain invariant under Γ . A necessary and sufficient condition is that they remain invariant under the generators of Γ , which are SLARs.

Thus, we search the eigenfunctions invariant under an arbitrary SLAR G . For a given value k , let us call their space $\mathcal{V}_G^k \subset \mathcal{V}^k$. Since G preserves the vector space $\mathcal{V}^{k, M}$, we have the decomposition

$$\mathcal{V}_G^k = \bigoplus_{M=-k/2}^{k/2} \mathcal{V}_G^{k, M}; \quad \mathcal{V}_G^{k, M} \equiv \mathcal{V}^{k, M} \cap \mathcal{V}_G^k.$$

The search of the G -invariant eigenmodes is equivalent to the search of the vector spaces $\mathcal{V}_G^{k, M}$, for the different values of k, M , in $\mathcal{V}^{k, M}$.

In $\mathcal{V}^{k, M}$, the expansion of a function f on the canonical basis,

$$f = \sum_{m=-k/2}^{k/2} f_{k; M m} T_{k; M, m},$$

describes it as the vector F with the $k+1$ components $F_m \equiv f_{k; M m}$. Similarly, we write for simplicity the matrix $M_{mn} \equiv D_{nm}^{k/2}(u_\ell^{-1})$. Then, equ.(24) takes the very simple form

$$F_m = \sum_n M_{mn} F_n. \quad (25)$$

This is an eigenvalue equation $MF = F$. Its solutions are the eigenvectors of the matrix M , with the eigenvalue 1.

We recall (see equ.8) that the Wigner function are the rotation matrix elements in the $(k+1)$ - dimensional representation $k/2$. Assuming $k = 2\ell$ even, the vector F represents also an harmonic function on the 2-sphere $\psi_V \equiv \sum_m F_m Y_{\ell m} \in \mathcal{V}^\ell$. Thus, (25) becomes

$$\psi_V = (R_g)^t \psi_V$$

(the subscript t holds for matrix transposition). This is the condition for the function f_V to be invariant under the $SO(3)$ rotation g^{-1} . The solution of this problem is well known. Let us call $\mathbf{n}$ and $2\pi/N$ the oriented (unit) axis and angle (assumed an integer divisor of 2π) of the rotation g .

- **Diagonal case:** When $\mathbf{n}$ is along Oz, the rotation is diagonalized : the $SU(2)$ matrix, $u = \text{diag}(e^{i\pi/N}, e^{-i\pi/N})$, $[R_g]_{mn} = \delta_{mn} e^{2i\pi m/N}$ and, using (4) and (14),

$$\begin{aligned} D_{m_2 m}^{k/2}(u^{-1}) &= \delta_{m m_2} (e^{-i\pi/N})^{m+m_2} P_{k/2-m}^{(m_2-m, m+m_2)}(1) \\ &= e^{-2i\pi m/N} \delta_{m m_2}. \end{aligned} \quad (26)$$

The eigenspace corresponding to the eigenvalue 1 is thus $\text{span}(e_{N\mu})_{-J_N \leq \mu \leq J_N}$, after defining $J_N \equiv [k/2N]$ ($[\cdot]$ means integer part), and where e_i is the i^{th} basis vector. In other words, the non zero components of an invariant vector have as index an integer multiple of N . The dimension of this space is $2J_N + 1$.

Coming back to $SO(4)$, this results implies immediately that

$$\mathcal{V}_G^{k,M} = \text{span}(T_{k;M,N\mu}), \quad -J_N \leq \mu \leq J_N.$$

- When $\mathbf{n}$ is not along Oz, we may diagonalize R_g as $g = \rho^{-1} D \rho$. The eigenvalue equation becomes

$$f_V = \rho^{-1} D \rho f_V.$$

This is the same eigenvalue equation as above, with the same solution, although not for V but for ρV . The eigenspace corresponding to the eigenvalue 1 is thus $\text{span}(\rho^{-1} e_{N\mu})_{-J_N \leq \mu \leq J_N}$.

Coming back to $SO(4)$, this results implies immediately that

$$\mathcal{V}_G^{k,M} = \text{span}(\rho^{-1} T_{k;M,N\mu}), \quad -J_N \leq \mu \leq J_N.$$

4 Eigenmodes of spherical spaces

When we have two generators $G_\pm$, the eigenspace is simply

$$\mathcal{V}_X^{k,M} = \mathcal{V}_{G_+}^{k,M} \cap \mathcal{V}_{G_-}^{k,M}.$$

In any spherical space, we chose a frame where one of the generators of Γ is diagonalized, so that the invariance condition relative to this generator takes the simple form (29). Thus, the problem is reduced to the search of the invariance condition relative to the other generators.

4.1 Diagonal matrices

Applying equation (23) to a diagonal matrix as above, we obtain:

$$R_g T_{k;m_1,m_2}(v) = e^{-2i\pi m_2/N} T_{k;m_1 m_2}(v) \quad (27)$$

Let us assume that a generator of the holonomy group of a spherical space X is represented by a diagonal left action $SU(2)$ matrix $g \approx (u_l, \mathbb{I})$ with u_l of the diagonal from above. The invariance condition, under g , of the basis functions for this space takes the form

$$T_{k;m_1,m_2}(u_l) = e^{-2i\pi m_2/N} T_{k;m_1 m_2}(u_l). \quad (28)$$

Its solution is

$$m_2 = 0 \pmod{N}, \quad (29)$$

or $m_2 = N\mu$, where μ varies in the range $[-J_N..J_N]$.

Hereafter, all sums involving μ, ν are assumed to go in this range $(2J_N + 1)$ values). This is the selection rule for the eigenmodes invariant under g .

4.2 A general procedure

The spherical spaces considered below all have two generators which are SLARs, $G_{\pm} \equiv (\gamma_{\pm}, \mathbb{I})$. Let us chose a frame where G_+ is diagonal. Then, form above, we can write

$$\mathcal{V}_G^{k,M} = \text{span}(T_{k;M,N\mu}), \quad -J_N \leq \mu \leq J_N.$$

In other words, all eigenfunctions $T_{k;M,m}$, with $m = N\mu$ are G_+ -invariant.

Since there is no simple way to calculate the intersection

$$\mathcal{V}_X^{k,M} = \mathcal{V}_{G_+}^{k,M} \cap \mathcal{V}_{G_-}^{k,M},$$

we propose a practical procedure.

Let us expand a G_+ -invariant function in the basis $(\mathcal{B}_2)$ as

$$f = \sum_m \sum_{\mu} f_{k;m,N\mu} T_{k;m,N\mu}, \quad \mu = -J_N \dots J_N.$$

The invariance condition under the rotation γ_- , for such a function, takes the form:

$$\begin{aligned} R_{G_-} f &= f : \\ \sum_m \sum_{\mu} f_{k;m,N\mu} R_{G_-} T_{k;m,N\mu}(x) &= \sum_m \sum_{\mu} f_{k;m,N\mu} T_{k;m,N\mu}(x). \end{aligned} \quad (30)$$

With the transformation law (23), this becomes:

$$f_{k;m,N\mu} = \sum_{\nu} f_{k;m,N\nu} D_{N\nu,N\mu}^{k/2} ((\gamma_-)^{-1}). \quad (31)$$

This equation is independant of index m .

This implies that the eigenspace for k splits as the direct sum

$$\mathcal{V}_X^k = \bigoplus_m \mathcal{V}_X^{k,m}.$$

The degeneracy of the eigenvalue λ_k is the dimension of $\mathcal{V}_X^k$. This implies that

$$\dim(\mathcal{V}_X^k) = (k+1) \dim(\mathcal{V}_X^{k,m}).$$

Thus, the solution for m does not depend on m , and we can write $f_{k;m,n} = f_{k;m_0,n}$, where m_0 is some index in $[-k/2; k/2]$. Equ. (31) is still an eigenvalue equation, but in a vector space of restricted dimension $2J_N + 1$: Let us define the matrix $\mathcal{M}_k$, of order $2J_N + 1$, through its coefficients:

$$[\mathcal{M}_k]_{\mu,\nu} \equiv D_{N\nu,N\mu}^{k/2} ((\gamma_-)^{-1}), \quad \mu, \nu = -J_N \dots J_N.$$

Note that this matrix does not depend on the index M . Let us also define the $(2J_N + 1)$ - vector $\mathcal{F}$ through its components :

$$[\mathcal{F}]_\mu \equiv f_{k;M,N\mu}.$$

Equation (31) could thus be written in matrix form:

$$\mathcal{M}_k \mathcal{F} = \mathcal{F}. \quad (32)$$

This eigenvalue equation may be easily solved, for instance with Mathematica. The results are given in appendix.

Having found the solution under the form of the vectors $\mathcal{F}_s$, with $1 \leq s \leq \sigma$, with $\sigma = 1/(2J_N + 1)$ times the degeneracy of the eigenvalue (σ is found as a result of the eigenvalue problem; the value coincide with that given by [6]). We have finally a basis $(\mathcal{Y}_{k;M,s})_{M=-k/2 \dots k/2, s=1 \dots \sigma}$ for the eigenmodes of X corresponding to the eigenvalue k , where

$$\mathcal{Y}_{k;M,s} \equiv \sum_{\mu=-J_N}^{J_N} (\mathcal{F}_s)_\mu T_{k;M,N\mu}. \quad (33)$$

5 Application to the spherical spaces

For each space, the steps are the following:

- We write the two generators in the $SU(2) \times SU(2)$ forms. Since both are SLARs, this corresponds to two $SU(2)$ matrices $\gamma_\pm$. (We thank J. Weeks for providing generators in suitable form).
- We diagonalize the matrix γ_+ .
- We deduce the value J_N .
- We solve the eigenvalue equation (32) in the vector space of dimension $2J_N + 1$. This gives the vector(s) $\mathcal{F}$ by their components
- Then equ. (33) gives the eigenmodes of X .

5.1 Dodecahedral space

The two $SU(2)$ generators,

$$\begin{pmatrix} c + iC & \pm i/2 \\ \pm i/2 & c - iC \end{pmatrix}; \quad c \equiv \cos(\pi/5), \quad C \equiv \cos(2\pi/5), \quad (34)$$

become, after diagonalization,

$$\gamma_+ = \begin{pmatrix} e^{i\pi/5} & 0 \\ 0 & e^{-i\pi/5} \end{pmatrix}$$

$$\gamma_- = \begin{pmatrix} \cos(\pi/5) + i \sin(\pi/5)/\sqrt{5} & 2i \sin(\pi/5)/\sqrt{5} \\ 2i \sin(\pi/5)/\sqrt{5} & \cos(\pi/5) - i \sin(\pi/5)/\sqrt{5} \end{pmatrix}. \quad (35)$$

We have $N = 5$, $J_N = \lfloor k/10 \rfloor$.

Equ.(33) gives the eigenmodes. We have checked that they correspond to those found by [9].

As we mentioned, it may be more convenient to express these modes in the more widely used basis $(\mathcal{B}_1)$. The solution of our problem in this basis is, for the dodecahedral space:

$$f_k(x) = \sum_{l,m,m'} i^{m-k-l} (-1)^{m'} f_{k;m'}(\frac{k}{2}, m' - m; \frac{k}{2}, m' \mid l, m) \mathcal{Y}_{klm}, \quad (36)$$

where the index m' satisfy the condition (29) for the dodecahedral space, l vary in $\llbracket 0, k \rrbracket$ and m vary in $\llbracket -l, l \rrbracket$.

5.2 Other multi-connected spherical spaces

Beside lens and prism spaces, three other multi-connected spherical spaces remain. In each one, holonomy groups are generated by two left-action $\text{SO}(4)$ rotations. We follow the same procedure.

We give in the table below a list of these spaces with their 2 generators, under their $\text{SU}(2)$ form.

Tetrahedron	$\frac{1}{2} \begin{pmatrix} 1-i & -1-i \\ 1+i & 1+i \end{pmatrix}$	$\frac{1}{2} \begin{pmatrix} 1+i & -1+i \\ 1+i & 1-i \end{pmatrix}$
diagonal form	$\begin{pmatrix} e^{i\pi/3} & 0 \\ 0 & e^{-i\pi/3} \end{pmatrix}$	$\frac{1}{\sqrt{3}} \begin{pmatrix} \frac{1}{2}(\sqrt{3}+i) & -1-i \\ 1-i & \frac{1}{2}(\sqrt{3}-i) \end{pmatrix}$

Octahedron	$\begin{pmatrix} e^{-i\pi/2} & 0 \\ 0 & e^{i\pi/2} \end{pmatrix}$	$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}$
------------	---	--

Icosahedron	$\begin{pmatrix} c+i/2 & iC \\ iC & c-i/2 \end{pmatrix}$	$\begin{pmatrix} c+iC & 1/2 \\ -1/2 & c-iC \end{pmatrix}$
diagonal form	$\begin{pmatrix} e^{i\pi/5} & 0 \\ 0 & e^{-i\pi/5} \end{pmatrix}$	$\begin{pmatrix} c+is & 1/2-ics \\ -1/2-ics & c-is \end{pmatrix}$

with $c := \cos(\pi/5)$ and $s := \sin(\pi/5)/\sqrt{5}$.

Computations gives the tables in appendix, where we presented the first modes for the different spherical spaces. More modes can be obtain by sending us an email at: cailleri@discovery.saclay cea.fr.

Note that the modes for the Icosahedron are the same than those of the Dodecahedron, as expected; they are however given here in a different frame.

6 Conclusion

The introduction of the basis $(\mathcal{B}_2)$ of Wigner functions provided the explicit behavior of the modes of $\mathcal{S}^3$, under the rotations of $\text{SO}(4)$. This allowed to write an invariance condition of these modes under any rotation, and under the holonomy group Γ of a spherical space $\mathcal{S}^3/\Gamma$. The solutions of the corresponding equations give the vector spaces of the eigenmodes of $\mathcal{S}^3/\Gamma$. These modes are calculated explicitly, for the first values of k , for the spherical spaces, in the basis $(\mathcal{B}_2)$. We give the transformation relations to express them in the more usual basis $(\mathcal{B}_1)$ of hyperspherical harmonics.

This allows to calculate the power angular spectrum of the CMB fluctuations, as predicted by the cosmological model with Poincaré dodecahedral space, up to arbitrary multipoles k . This work is in progress. It will allow to check if the predictions of [13] hold up to higher orders. In any case, the confrontation of the predicted power spectrum with observations may not offer enough discriminating power. Thus, we intend to calculate the cross correlations predicted from this model. They are zero for Gaussian isotropic fluctuations, and thus allow more specific tests of cosmic topology. This work is also in progress.

The present results allow to extend these calculations to the remaining spherical spaces, a work also in progress. Hopefully, this will provide a definitive test of the case of a multi-connected space with positive curvature, at least for a range of scales comparable with that of the last scattering surface.

References

- [1] *Group Theory and the Hydrogen Atom (I)*, Bander M. & Itzykson C., Rev. Mod. Phys. 18,2,1966
- [2] *Angular momentum on quantum mechanics*, Edmonds, Princeton University Press 1960
- [3] *Higher transcendental Functions* A. Erdélyi, W. Magnus, F. Oberhettinger, F. G. Tricomi, McGraw-Hill 1953
- [4] A. Fryant, SIAM J. Math. Anal., Vol. 22, N 1, pp. 268-271, 1991
- [5] Gazeau J-P 1978, Ph.D. Thesis (Doctorat d'État ès Sciences Physiques Université Paris 6), *Nature algébrique de l'opérateur sturmien associé à l'équation de Schrödinger. Extension de certains résultats à la théorie des représentations finies de $GL(n, \mathbb{C})$ et de $U(n)$ (polynômes à treillis de Gel'fand)*.
- [6] Ikeda A. 1995, Kodai Math. J. 18 (1995) 57-67
- [7] Lachièze-Rey M. 2003, Journal of Physics A: Mathematical and General, volume 37, issue 1, pages 205 - 210 (<http://fr.arxiv.org/abs/math.SP/0304409>)
- [8] Lachièze-Rey M. 2004, Journal of Physics A: Mathematical and general, vol. 37, 21, p 5625-5634 (<http://fr.arxiv.org/abs/math.SP/0401153>)
- [9] Lachièze-Rey M. 2004, C.Q.G. 21 (2004) 1-10 (gr-qc/0402035)
- [10] Lachièze-Rey M., Luminet J.-P. 1995, Phys.Rept. 254 (1995) 135-214
- [11] Lehoucq R., Uzan J.-P., Weeks J. 2002, Kodai math. J. 26 (2003), 119-136 (math.SP/0202072)
- [12] Lehoucq R., Weeks J., Uzan J. P., Gausmann E., Luminet J.-P., Class.Quant.Grav. 19 (2002) 4683-4708 (gr-qc/0205009)
- [13] Luminet J.-P. Weeks J. R., Riazuelo A., Lehoucq R. & Uzan J.-P. 2003, Nature 425 (9 october 2003), p.593
- [14] *Fonctions spéciales de la physique mathématique*, A. Nikiforov, V. Ouvarov, Editions Mir (Moscou) 1983
- [15] A. Riazuelo, J.-P. Uzan, R. Lehoucq, J. Weeks, Phys.Rev. D69 (2004) 103514 (astro-ph/0212223)

Table of eigenfunctions for the dodecahedral space

Eigenmodes of dodecahedral space corresponds to the basis of functions $(f_k(x))_{m \in \llbracket -k/2; k/2 \rrbracket}$, where $f_k = \sum_{m, m'} f_{k; m'} T_{k; m, m'}$. In this table is indicated the allowed k (those where D_k admits 1 as an eigenvalue) and the corresponding $f_{k; m'}$, for $m' = 0 \pmod{5}$ (for others values of m' , $f_{k; m'} = 0$).

k=12			
0.529150262 i	0.663324958	0.529150262 i	
k=20			
-0.3158058475 i	0.578273291	0.362950869 i	0.578273291
-0.3158058475 i			
k=24			
-0.486949689	0.3025227264 i	0.585422924	0.3025227264 i
-0.486949689			
k=30			
-0.2829840984	0.391305507 i	-0.516526862	0
0.516526862	-0.391305507 i	0.2829840984	
k=32			
-0.329930901 i	0.425449096	-0.331083071 i	0.448380790
-0.331083071 i	0.425449096	-0.329930901 i	

For this space and others, we have calculated more modes, and with an higher precision. As indicated in the text, these modes could be asked at: cailleri@discovery.saclay.cea.fr.

Table of eigenfunctions for the tetrahedral space

For this space, we present modes under the form: $f_{k;m}$, with the notation used above, where $m = 0 \pmod{3}$. For $k = 12$, we can observe that modes are degenerated and we thus have 2 series of $f_{k;m}$.

k=6	
0.333333333 - 0.333333333 i	0.745355992
-0.333333333 - 0.333333333 i	
k=8	
0.608580619	-0.360041149 - 0.360041149 i
-0.608580619 i	
k=12	
-0.00323332162 + 0.00323332162 i	0.738332205
0.2666917602 + 0.2666917602 i	0.2828693156 i
-0.341013640 + 0.341013640 i	
0.354280119 + 0.354280119 i	-0.0861446455 i
-0.353087976 + 0.353087976 i	0.689157164
-0.0929259330 - 0.0929259330 i	
k=14	
0.491351820 i	-0.1888525745 + 0.1888525745 i
0.611952283	0.1888525745 + 0.1888525745 i
-0.491351820 i	

Table of eigenfunctions for the octahedral space

For this space, we present modes under the form: $f_{k;m}$, with the notation used above, where $m = 0 \pmod{2}$.

k=4			
-0.612372435	0.500000000	-0.612372435	
k=8			
-0.379649579	-0.1447862189	-0.818416944	-0.1447862189
-0.379649579			
-0.530330085	0.3118047822	-0.493006648	0.3118047822
-0.530330085			
k=10			
-0.353553390	-0.612372435	0	0.612372435
0.353553390			
k=12			
0.507752400	-0.1250000000	0.342326598	-0.467707173
0.342326598	-0.1250000000	0.507752400	
0.1230304012	-0.672015232	0.0829470795	0.2296903537
0.0829470795	-0.672015232	0.1230304012	

Table of eigenfunctions for the icosahedral space

For this space, we present modes under the form: $f_{k;m}$, with the notation used above, where $m = 0 \pmod{5}$.

k=12	
0.1882530268 + 0.3999418963 i	0.6533724434
-0.2617387416 + 0.5560616500 i	
k=20	
-0.2115446757 - 0.0542397647 i	0.3004753102 - 0.3633777752 i
0.1486544684 + 0.3158151081 i	0.655649414
0.1798126817 - 0.3820104577 i	
k=24	
-0.0622550053 - 0.3255605557 i	-0.2352778614 - 0.0603249209 i
-0.3529784282 + 0.4268720642 i	-0.1437372075 - 0.3053684305 i
0.6406372434	
k=30	
0.02978235622 + 0.1557458776 i	0.2506842548 + 0.0642750991 i
-0.2564612869 + 0.3101497151 i	0.0002877519 + 0.0006113259 i
0.560360227	0.2128816958 - 0.4522652865 i
0.2718985718 + 0.3288186907 i	